

Multidimensional Cosmology: Spatially Homogeneous models of dimension $4+1$

Sigbjørn Hervik*

DAMTP,
Centre for Mathematical Sciences,
Cambridge University,
Wilberforce Rd.,
Cambridge CB3 0WA, UK

February 7, 2008

Abstract

In this paper we classify all $4+1$ cosmological models where the spatial hypersurfaces are connected and simply connected homogeneous Riemannian manifolds. These models come in two categories, multiply transitive and simply transitive models. There are in all five different multiply transitive models which cannot be considered as a special case of a simply transitive model. The classification of simply transitive models, relies heavily upon the classification of the four dimensional (real) Lie algebras. For the orthogonal case, we derive all the equations of motion and give some examples of exact solutions. Also the problem of how these models can be compactified in context with the Kaluza-Klein mechanism, is addressed.

1 Introduction

The idea that our world has more dimensions than three is actually older than the theory of general relativity itself. Already in 1914, G. Nordström¹ put forward a five-dimensional scalar-tensor theory in an effort to unify gravity and electromagnetism. Since it was based upon his own theory of gravitation which was soon superseded by Einstein's theory, this work was neglected for several decades.

However, in 1919, T. Kaluza [2] constructed a similar unified theory of gravity and electromagnetism based on the linearized version of the general theory of relativity. Kaluza's work, which was published in 1921 and was followed by two important papers by Klein [3, 4], had a very interesting result: five-dimensional

*S.Hervik@damtp.cam.ac.uk

¹English translations of this and other related works are found in the book [1].

Einstein gravity could be seen upon as Einstein gravity in four dimensions plus electromagnetism.

The original idea had only one extra dimension, but string theorists today believe that we can have up to seven extra dimensions[5]. Supposedly, six of these have to be curled up in a small Calabi-Yau manifold[6]. Hence, among theorists the question is not *if* we have extra dimensions, but rather “*How many extra dimensions are there?*” and “*What is their nature?*”

In this paper we will consider the simplest models, models with one extra dimension. We will investigate cosmological models of 4+1 dimensions. In 3+1 dimensions the classification relies on Bianchi’s classification of the homogeneous three-manifolds [7]. Bianchi’s work was later generalised to the four-dimensional manifolds by Fubini [8]. In three dimensions the homogeneous manifolds have a special role. According to the Thurston conjecture, the homogeneous manifolds of dimension three are intimately related to the classification of three-manifolds[9, 10]. There does not exist a similar conjecture in four dimensions; on the contrary, topology in four dimensions is completely different. Notwithstanding, we will assume that our 4+1 cosmological model is *spatially homogeneous*. This assumption heavily restricts the number of possible four-manifolds to only a finite and manageable number. We know that our universe is homogeneous on scales larger than a billion light-years, and thus the assumption of homogeneity is by no means a radical and unrealistic one².

The purpose of our study is to investigate the impact of extra dimensions on the cosmic evolution of our universe. This may give us some understanding of how extra dimensions generically influence the observed 3+1 dimensional universe.

The simplest extension we can think of, going from three to four spatial dimensions, is just to assume that we have have a product space:

$$\Sigma \cong M \times S^1 \tag{1}$$

where Σ is a four dimensional spatial manifold and M a three dimensional manifold. However, this naive assumption does not necessarily need to be true. Experimental data from particle accelerators indicate that the size of this small extra dimension must be less than 10^{-18}m . Hence, since the size of the base-space M is greater than billions of light-years, the extra dimension may also be “twisted”. This twisting is at a global scale, and will therefore be unmeasurable in particle accelerators. However, we know that once in the past, the size of the universe might have been comparable with the small extra dimensions. Actually, it is one of our principal aims to try to explain why one (or more) dimension(s) is so incredibly much smaller than the three large ones we see today.

Some work along similar lines already exist in the literature. Forgacs and Horvath [11, 12] investigated already in 1979 how higher dimensional models could influence the cosmology of isotropic FRW universes. This work was followed by Chodos and Detweiler[13] who studied the translational invariant higher-dimensional cosmological models – also over 20 years ago. Later, other people have investigated other models [14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28]. Also, Lorenz-Petzold produced a string of papers

²From CMB measurements we know that the fluctuations away from homogeneity are no larger than 10^{-5} at the last scattering surface.

[29, 30, 31, 32, 33, 34, 35, 36] by lifting the homogeneous models in 3+1 dimensions, one by one, to higher dimensions. Maybe the closest work related to this (at least to the authors knowledge) are two articles in the mid-eighties [37, 38]. However, there does not seem to be any work which tries to classify all the spatially homogeneous spacetimes in 4+1 dimensions and gives a systematic approach to the equations of motion. The aim of this paper is exactly to do this; give a classification and with the aid of the orthonormal frame formalism, we will derive all the equations of motion.

The paper is organised as follows. First we classify all the multiply transitive spaces. We will throughout our paper assume, unless stated otherwise, that our space is connected and simply connected. It is of special interest to find all the homogeneous spaces which cannot be considered as simply transitive spaces. The simply transitive spaces are classified in section 3. This classification heavily relies upon the classification of the four-dimensional Lie algebras. We write down all the equations of motion and go on and provide with some examples and give some exact solutions. Lastly, we go to the question of compactification which is one of the key ingredients of the Kaluza-Klein mechanism.

2 Multiply transitive models

Let us first classify all the multiply transitive models which are connected and simply connected. In dimension three there is only one multiply transitive model which cannot be considered as a special case of a simply transitive one. This is the well-known Kantowski-Sachs model. The symmetry group in this case is four-dimensional and it has three-dimensional subgroup which acts on a two-sphere S^2 .

Our analysis is based on the classification of the homogeneous Riemannian spaces of dimension four due to Ishihara [39]³. Large parts of our results in this section can be extracted directly from this paper. In four dimensions there are in all 5 different models which cannot be seen as special cases of a simply transitive space. In the following, these will be emphasized.⁴

2.1 Maximally symmetric: $\dim \text{Isom}(\Sigma) = 10$

There are three (orientable) maximally symmetric spaces of dimension 4. They are the well-known cases: $\underline{S^4}$, $\mathbb{E}^4$ and $\mathbb{H}^4$. These have an isotropy group isomorphic to $SO(4)$. They correspond to the three different FRW cosmological models in 5D. Both the Euclidean space and the hyperbolic space have subgroups of the isometry group which acts simply transitive on their respective spaces. $\underline{S^4}$ on the other hand, has no proper subgroup that acts transitively on the space at all. Hence, $\underline{S^4}$ cannot be considered as a special case of a simply transitive space.

³The homogeneous Riemannian spaces are treated in a more general way in for example [40]. However, dimension four seems to be a special case. This is mainly due to the non-simpleness of $SO(4)$.

⁴The underlined spaces will correspond to these multiply transitive models. However, they will only be underlined in the section were they have the smallest transitive symmetry group.

2.2 Kähler manifolds: $\dim \text{Isom}(\Sigma) = 8$

Interestingly, there also exist four-dimensional Riemannian spaces which have an 8-dimensional isometry group⁵. These are the three Kähler manifolds of constant curvature: $\mathbb{CP}^2$, $\mathbb{C}^2$ and⁶ $\mathbb{H}_{\mathbb{C}}^2$. These spaces have an isotropy subgroup isomorphic to $U(2)$. Both $\mathbb{C}^2$ and $\mathbb{H}_{\mathbb{C}}^2$ have a simply transitive subgroup.

The Riemann curvature tensor for these spaces can be written in complex coordinates as

$$R_{\alpha\bar{\beta}\delta\bar{\gamma}} = K (g_{\alpha\bar{\beta}}g_{\delta\bar{\gamma}} + g_{\alpha\bar{\gamma}}g_{\delta\bar{\beta}}) \quad (2)$$

where $K > 0$, $K = 0$ and $K < 0$ for $\mathbb{CP}^2$, $\mathbb{C}^2$ and $\mathbb{H}_{\mathbb{C}}^2$ respectively. Thus these spaces have constant holomorphic sectional curvature. The isotropy group is the group of motions that leaves the Kählerian structure invariant.

2.3 $\dim \text{Isom}(\Sigma) = 7$

Spaces that possess a 7-dimensional isometry group come in three classes. The first class consists of maximally symmetric three-manifolds times a line: $\mathbb{E}^1 \times S^3$, $\mathbb{E}^4$ and $\mathbb{E}^1 \times \mathbb{H}^3$. These have an isotropy group isomorphic to $SO(3)$. The second category consists of $\mathbb{H}^4$ with a certain 7-dimensional symmetry group. The third is the Kähler manifold $\mathbb{C}^2$ with isotropy group $SU(2)$. All of these spaces can be considered as a special case of a simply transitive space.

2.4 $\dim \text{Isom}(\Sigma) = 6$

All of these spaces are products of two two-dimensional maximally symmetric spaces: $S^2 \times S^2$, $S^2 \times \mathbb{E}^2$, $S^2 \times \mathbb{H}^2$, $\mathbb{H}^2 \times \mathbb{E}^2$, $\mathbb{H}^2 \times \mathbb{H}^2$ and $\mathbb{E}^4$. These have the isotropy group $SO(2) \times SO(2)$. Only $S^2 \times S^2$ does not have a lower dimensional subgroup of the isometry group which acts transitively on the spatial hypersurface. For this model, the metric can be written as

$$ds^2 = -dt^2 + a(t)^2 (d\theta^2 + \sin^2 \theta d\phi^2) + b(t)^2 (d\psi^2 + \sin^2 \psi d\xi^2). \quad (3)$$

2.5 $\dim \text{Isom}(\Sigma) = 5$

There are several spaces possessing a 5-dimensional isometry group. Only two of them cannot be considered as a special case of a simply transitive space: $S^2 \times \mathbb{E}^2$ and $S^2 \times \mathbb{H}^2$. The isotropy group for both these spaces is $SO(2)$. Examples of such models for the case $S^2 \times \mathbb{E}^2$ case are

$$ds^2 = -dt^2 + a(t)^2 (d\theta^2 + \sin^2 \theta d\phi^2) + b(t)^2 (e^{-2\beta(t)} dx^2 + e^{2\beta(t)} dy^2), \quad (4)$$

and in the case $S^2 \times \mathbb{H}^2$

$$ds^2 = -dt^2 + a(t)^2 (d\theta^2 + \sin^2 \theta d\phi^2) + b(t)^2 (e^{-2\beta(t)} dx^2 + e^{2\beta(t)+2Kx} dy^2). \quad (5)$$

The functions $a(t)$, $b(t)$ and $\beta(t)$ will be determined by the field equations.

⁵In general for a homogeneous Riemannian space M of dimension $n \neq 4$ there does not exist a closed subgroup of $\text{Isom}(M)$ of dimension r such that $n(n+1)/2 > r > 1 + n(n-1)/2$. These are the counterexamples for $n = 4$.

⁶For those who are unfamiliar with the complex hyperbolic spaces, consult for example Goldman's book [41].

3 Simply transitive models

In the simply transitive case, we can systematically construct the homogeneous spaces using the classification of the 4-dimensional Lie algebras. For a given simply transitive space, the Killing vectors obey a certain commutator relation

$$[\xi_i, \xi_j] = \tilde{C}_{ij}^k \xi_k \quad (6)$$

where the structure constants $\tilde{C}_{ij}^k$ are functions of t only. In the following we will follow Ellis and MacCallum [42] closely. However we will use a slightly different notation: Greek indices $(\mu, \nu, \dots)$ have range 0-4 over the full five dimensional space-time; lower case Latin indices $(a, b, i, j, \dots)$ have range 1-4 over the four-dimensional spatial hypersurfaces; upper case Latin indices $(A, B, \dots)$ have range 1-3 over three spatially directions. We will assume that the model is orthogonal, i.e. we assume that our spacetime is of the form

$$\Sigma_t \times \mathbb{R} \quad (7)$$

where the fluid four-velocity $\mathbf{u}$ is orthogonal to Σ_t .

We can now introduce a left-invariant spatial frame $\mathbf{e}_a$, which together with the fluid four-velocity $\mathbf{u} = \mathbf{e}_t$ forms a orthonormal frame denoted by $\mathbf{e}_\mu$. These commute, by definition, with the Killing vectors:

$$[\xi_i, \mathbf{e}_\mu] = 0. \quad (8)$$

The fünfbein $\mathbf{e}_\mu$ will now obey the commutation relations

$$[\mathbf{e}_\mu, \mathbf{e}_\nu] = \gamma_{\mu\nu}^\rho \mathbf{e}_\rho. \quad (9)$$

We can relate these commutator function to the connection coefficients for the particular orthonormal basis. For an orthonormal basis we define the rotation forms by

$$d\mathbf{e}_\mu = \mathbf{e}_\nu \otimes \Omega_{\mu}^\nu. \quad (10)$$

Denoting the dual frame of $\mathbf{e}_\mu$ by ω^μ , the rotation forms can be related to the connection coefficients via

$$\Omega_{\mu}^\nu = \Gamma_{\mu\lambda}^\nu \omega^\lambda. \quad (11)$$

Also, they possess the antisymmetry $\Omega_{\mu\nu} = -\Omega_{\nu\mu}$. Hence, the connection coefficients $\Gamma_{\mu\nu}^\rho$ for the orthonormal frame possess the antisymmetry

$$\Gamma_{\rho\mu\nu} = -\Gamma_{\mu\rho\nu} \quad (12)$$

and can be written as

$$\begin{aligned} \gamma_{\mu\nu}^\rho &= -(\Gamma_{\mu\nu}^\rho - \Gamma_{\nu\mu}^\rho) \\ \Gamma_{\rho\mu\nu} &= -\frac{1}{2}(\gamma_{\rho\mu\nu} + \gamma_{\mu\nu\rho} - \gamma_{\nu\rho\mu}). \end{aligned} \quad (13)$$

The orthogonality condition requires that

$$\gamma_{0a}^0 = \gamma_{ab}^0 = 0. \quad (14)$$

We can split the remaining part of the structure constants into

$$\begin{aligned}\gamma^a_{0b} &= -\theta^a_b + \Omega^a_b \\ \gamma^i_{jk} &= C^i_{jk}.\end{aligned}\tag{15}$$

Here are Ω_{ab} the angular velocity in the ab -plane of a Fermi-propagated axis with respect to the triad $\mathbf{e}_a$, and θ^a_b the volume expansion tensor. If u^μ is the time-like vector-field orthogonal to the hypersurfaces Σ_t , then θ^a_b is defined by $\theta_{\mu\nu} = u_{\mu;\nu}$. One can readily see that $\theta_{\mu\nu}$ is symmetric and $\theta_{\mu\nu}u^\mu = 0$.

We can further split the volume expansion tensor into a trace and trace-free part

$$\theta_{\mu\nu} = \frac{1}{4}h_{\mu\nu}\theta + \sigma_{\mu\nu}.\tag{16}$$

The tensor $\sigma_{\mu\nu}$ is the shear tensor and $h_{\mu\nu} = g_{\mu\nu} - u_\mu u_\nu$ is the projection tensor onto the hypersurfaces Σ_t .

The structure constants C^k_{ij} defines the specific Lie algebra under consideration. These can be separated into a trace (or vector) part and a trace-free part

$$C^k_{ij} = D^k_{ij} + \delta^k_i a_j - \delta^k_j a_i\tag{17}$$

where

$$\begin{aligned}a_i &= \frac{1}{3}\text{Tr}(C^k_{ji}) \equiv \frac{1}{3}C^j_{ji} \\ D^k_{ij} &= C^k_{ij} - \frac{2}{3}C^l_{[i|l]}\delta^k_{j]}.\end{aligned}\tag{18}$$

The Lie algebra given by the structure constants C^k_{ij} is isomorphic to the Lie algebra $\tilde{C}^k_{ij}$ defined in eq. (6).

The Jacobi identity,

$$[\mathbf{e}_\mu, [\mathbf{e}_\nu, \mathbf{e}_\rho]] + [\mathbf{e}_\nu, [\mathbf{e}_\rho, \mathbf{e}_\mu]] + [\mathbf{e}_\rho, [\mathbf{e}_\mu, \mathbf{e}_\nu]] = 0\tag{19}$$

reduces now to the following constraint equations

$$a_k D^k_{ij} = 0\tag{20}$$

$$D^d_{ab}D^e_{cd} + D^d_{ca}D^e_{bd} + D^d_{bc}D^e_{ad} + D^e_{ab}a_c + D^e_{ca}a_b + D^e_{bc}a_a = 0\tag{21}$$

and evolution equations

$$\dot{a}_b + \frac{1}{4}\theta a_b - (\Omega^d_b - \sigma^d_b)a_d = 0\tag{22}$$

$$\dot{D}^a_{bc} + \frac{1}{4}\theta D^a_{bc} + 2\left(\Omega^d_{[b} - \sigma^d_{[b}\right)D^a_{c]d} + (\Omega^a_d - \sigma^a_d)D^d_{bc} = 0.\tag{23}$$

Equation (21) is just the Jacobi identity for C^k_{ij} in disguise, while eq. (20) is the trace of eq. (21).

3.1 The Field Equations

Let us assume that the energy-momentum tensor is of the form

$$T_{\mu\nu} = (\rho + p)u_\mu u_\nu + pg_{\mu\nu} + \pi_{\mu\nu} \quad (24)$$

where $\pi_{\mu\nu}$ is the anisotropic stress tensor. This tensor is symmetric and has the properties

$$\pi^\mu{}_\mu = u^\mu \pi_{\mu\nu} = 0. \quad (25)$$

The identity $T^\mu{}_{\nu;\mu} = 0$ leads to the energy conservation equation

$$\dot{\rho} + (\rho + p)\theta + \pi^{\mu\nu}\sigma_{\mu\nu} = 0. \quad (26)$$

Together with an equation of state for the fluid, these equations govern the evolution of the fluid in our model.

The Riemann curvature tensor is given by

$$R^\alpha{}_{\beta\gamma\delta} = \mathbf{e}_\gamma(\Gamma^\alpha{}_{\beta\delta}) - \mathbf{e}_\delta(\Gamma^\alpha{}_{\beta\gamma}) + \Gamma^\alpha{}_{\lambda\gamma}\Gamma^\lambda{}_{\beta\delta} - \Gamma^\alpha{}_{\lambda\delta}\Gamma^\lambda{}_{\beta\gamma} - \Gamma^\alpha{}_{\beta\lambda}\gamma^\lambda{}_{\gamma\delta}. \quad (27)$$

From this expression we can readily calculate the field equations. The 5D Raychaudhuri's equation is

$$\dot{\theta} + \frac{1}{4}\theta^2 + \sigma^{ab}\sigma_{ab} + \frac{2}{3}(\rho + 2p - \Lambda) = 0 \quad (28)$$

while the $(0, a)$ equations are

$$4\sigma_{ac}a^c - \sigma^b{}_c D^c{}_{ba} = 0. \quad (29)$$

The trace of the field equations yields the 5D Friedmann equation (or the constraint equation):

$$\begin{aligned} \frac{3}{8}\theta^2 &= \frac{1}{2}\sigma^{ab}\sigma_{ab} + 6a^2 + \frac{1}{8}\mathfrak{D}^2 + \rho + \Lambda \\ \mathfrak{D}^2 &= D^a{}_{bc}D_a{}^{bc} + 2D^a{}_{bc}D^b{}_a{}^c. \end{aligned} \quad (30)$$

The shear equations can be derived from the trace-free part of the field equations:

$$\dot{\sigma}_{ab} + \theta\sigma_{ab} + \frac{3}{2}(D_{abd} + D_{bad})a^d + 2\sigma^c{}_{(a}\Omega_{b)c} + b_{ab} - \frac{1}{4}h_{ab}b^c{}_c = \pi_{ab} \quad (31)$$

where we have set

$$b_{ab} \equiv -\frac{1}{4}(2D^c{}_{ad}D^d{}_{cb} - D_{acd}D_b{}^{cd} + 2D^c{}_{ad}D^d{}_{bc}). \quad (32)$$

Note that $4b^a{}_a = -\mathfrak{D}^2$. The tensor b_{ab} can be interpreted as part of the spatial Ricci tensor:

$${}^{(4)}R_{ab} = b_{ab} + \frac{3}{2}(D_{abd} + D_{bad})a^d - 3h_{ab}a^2. \quad (33)$$

Hence, the curvature of the spatial four-surfaces is

$${}^{(4)}R = b^a{}_a - 12a^2 = -\frac{1}{4}\mathfrak{D}^2 - 12a^2. \quad (34)$$

3.2 The four-dimensional Lie algebras

In a series of papers Mubarakzyanov [43, 44, 45] gave a classification of real Lie algebras up to dimension 5.⁷ A list of all the algebras is given in [47, 48] which is more accessible for non-Russian readers (see also [49] for the four-dimensional case). Here we will be interested in the four-dimensional classification and we will be using mostly the notation in [47]. A useful reference is also a report by MacCallum [50].

We will investigate Lie algebras from two different point of views; one from a geometric point of view, and the other from an algebraic point of view. The two different ways of investigating Lie algebras both have their strengths and weaknesses and are useful for different purposes. We will start out from the geometrical point of view; dividing them into decomposable and indecomposable ones. The geometrical picture is completely different in these two different classes. The decomposable ones naturally give rise to product spaces⁸, while the indecomposable ones cannot be written as a topological product.

3.2.1 Decomposable Lie algebras

Since the three-dimensional Lie algebras are used quite frequently in the literature we will use a notation similar to the Bianchi types when the algebras are composed of these. The decomposable four-dimensional Lie algebras are

$$A_3 \oplus \mathbb{R}, \quad \text{and} \quad A_{2,1} \oplus A_{2,1} \quad (35)$$

where A_3 is one of the Bianchi type Lie algebras, labelled I-IX, and $A_{2,1}$ is the only non-trivial two-dimensional Lie algebra. In this notation, the extension of the type IX Bianchi type will be denoted $\text{IX} \oplus \mathbb{R}$. $A_{2,1}$ can be represented by the single non-trivial commutator

$$[\mathbf{e}_1, \mathbf{e}_2] = \mathbf{e}_1. \quad (36)$$

One can show that this Lie algebra acts simply transitively on $\mathbb{H}^2$. It has been known for a while that the commutators for the Bianchi Lie algebras can be written

$$C^A_{BC} = \epsilon_{BCD} n^{DA} + \delta^A_B a_C - \delta^A_C a_B \quad (37)$$

where n^{AD} is a symmetric matrix. This is called the *Behr decomposition* [51]. The Jacobi identity reduces to the single relation

$$n^{AD} a_D = 0. \quad (38)$$

The Bianchi types can now be classified in terms of the eigenvalues of the matrix n^{AD} [42].

3.2.2 Indecomposable Lie algebras

The classification of the indecomposable four-dimensional Lie algebras are listed in table 1.

⁷The classification of the *complex* Lie algebras up to dimension 6 was actually done by Sophus Lie himself already in the 1890's [46].

⁸As Lie groups.

Name	Commutator relations	a_i
$A_{4,1}$	$[\mathbf{e}_2, \mathbf{e}_4] = \mathbf{e}_1, [\mathbf{e}_3, \mathbf{e}_4] = \mathbf{e}_2$	0
$A_{4,2}^p$	$[\mathbf{e}_1, \mathbf{e}_4] = p\mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = \mathbf{e}_2,$ $[\mathbf{e}_3, \mathbf{e}_4] = \mathbf{e}_2 + \mathbf{e}_3 \quad (p \neq 0)$	$\frac{1}{3}(p+2)\delta_i^4$
$A_{4,3}$	$[\mathbf{e}_1, \mathbf{e}_4] = \mathbf{e}_1, [\mathbf{e}_3, \mathbf{e}_4] = \mathbf{e}_2$	$\frac{1}{3}\delta_i^4$
$A_{4,4}$	$[\mathbf{e}_1, \mathbf{e}_4] = \mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = \mathbf{e}_1 + \mathbf{e}_2,$ $[\mathbf{e}_3, \mathbf{e}_4] = \mathbf{e}_2 + \mathbf{e}_3$	δ_i^4
$A_{4,5}^{pq}$	$[\mathbf{e}_1, \mathbf{e}_4] = \mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = p\mathbf{e}_2, [\mathbf{e}_3, \mathbf{e}_4] = q\mathbf{e}_3$ $(pq \neq 0, -1 \leq q \leq p \leq 1)$	$\frac{1}{3}(1+p+q)\delta_i^4$
$A_{4,6}^{pq}$	$[\mathbf{e}_1, \mathbf{e}_4] = p\mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = q\mathbf{e}_2 - \mathbf{e}_3,$ $[\mathbf{e}_3, \mathbf{e}_4] = \mathbf{e}_2 + q\mathbf{e}_3 \quad (p \neq 0, q \geq 0)$	$\frac{1}{3}(p+2q)\delta_i^4$
$A_{4,7}$	$[\mathbf{e}_2, \mathbf{e}_3] = \mathbf{e}_1, [\mathbf{e}_1, \mathbf{e}_4] = 2\mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = \mathbf{e}_2,$ $[\mathbf{e}_3, \mathbf{e}_4] = \mathbf{e}_2 + \mathbf{e}_3$	$\frac{4}{3}\delta_i^4$
$A_{4,8}$	$[\mathbf{e}_2, \mathbf{e}_3] = \mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = \mathbf{e}_2, [\mathbf{e}_3, \mathbf{e}_4] = -\mathbf{e}_3$	0
$A_{4,9}^q$	$[\mathbf{e}_2, \mathbf{e}_3] = \mathbf{e}_1, [\mathbf{e}_1, \mathbf{e}_4] = (1+q)\mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = \mathbf{e}_2,$ $[\mathbf{e}_3, \mathbf{e}_4] = q\mathbf{e}_3 \quad (-1 < q \leq 1)$	$\frac{2}{3}(1+q)\delta_i^4$
$A_{4,10}$	$[\mathbf{e}_2, \mathbf{e}_3] = \mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = -\mathbf{e}_3, [\mathbf{e}_3, \mathbf{e}_4] = \mathbf{e}_2$	0
$A_{4,11}^q$	$[\mathbf{e}_2, \mathbf{e}_3] = \mathbf{e}_1, [\mathbf{e}_1, \mathbf{e}_4] = 2q\mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = q\mathbf{e}_2 - \mathbf{e}_3,$ $[\mathbf{e}_3, \mathbf{e}_4] = \mathbf{e}_2 + q\mathbf{e}_3, \quad (q > 0)$	$\frac{4}{3}q\delta_i^4$
$A_{4,12}$	$[\mathbf{e}_1, \mathbf{e}_4] = \mathbf{e}_1, [\mathbf{e}_2, \mathbf{e}_4] = \mathbf{e}_2,$ $[\mathbf{e}_1, \mathbf{e}_3] = -\mathbf{e}_2, [\mathbf{e}_2, \mathbf{e}_3] = -\mathbf{e}_1$	$\frac{2}{3}\delta_i^4$

Table 1: The indecomposable four-dimensional algebras

Note that some of the parametric limits yields various other Lie algebras. We have

$$\begin{aligned}
\lim_{p \rightarrow 0} A_{4,2}^p &= IV \oplus \mathbb{R}, & \lim_{q \rightarrow 0} A_{4,5}^{pq} &= VI_h \oplus \mathbb{R}, \\
\lim_{p \rightarrow 0} A_{4,6}^{pq} &= VII_h \oplus \mathbb{R}, & \lim_{q \rightarrow 0} A_{4,9}^q &= A_{4,8}, \\
\lim_{q \rightarrow 0} A_{4,11}^q &= A_{4,10}.
\end{aligned} \tag{39}$$

Unfortunately there does not exist a simple expression, as in the three-dimensional case, for the structure constants. This is perhaps the greatest obstacle to our analysis. However, in several cases, the general form of the structure constants can be worked out.

Let C_{ij}^k be a special representation of a Lie algebra $\mathcal{A}$. We define the space $\mathcal{W}(\mathcal{A})$ as

$$\mathcal{W}(\mathcal{A}) = \left\{ \tilde{C}_{ij}^k \mid \tilde{C}_{ij}^k = (\mathbf{A}^{-1})_l^k C_{nm}^l \mathbf{A}_i^n \mathbf{A}_j^m, \quad \mathbf{A} \in GL(4, \mathbb{R}) \right\}. \tag{40}$$

Two elements in $\mathcal{W}(\mathcal{A})$ correspond to two isomorphic Lie algebras. It would be convenient to have a specific parametrisation of $\mathcal{W}(\mathcal{A})$ for all of the four-dimensional Lie algebras. Unfortunately, such a parametrisation has not been found in the four-dimensional case. The union of $\mathcal{W}(\mathcal{A})$ over all possible Lie algebras in n -dimensions is called the *variety of n -dimensional Lie algebras*⁹. In dimensions two and three, the structure of this variety is known, but in four dimensions and higher it is not known in detail.

⁹For the case of complex Lie algebras of low dimensions, see [52].

However, for specific cases, $\mathcal{W}(\mathcal{A})$ can be found explicitly. For instance, for the pure vector type algebra, $A_{4,5}^{1,1}$ the structure constants are (see Table 1)

$$C_{ij}^k = \delta_i^k \delta_j^4 - \delta_j^k \delta_i^4. \quad (41)$$

Hence, $\mathcal{W}(A_{4,5}^{1,1})$ is the space of all non-zero vectors in $\mathbb{R}^4$:

$$\mathcal{W}(A_{4,5}^{1,1}) \cong \mathbb{R}^4 \setminus \{0\}. \quad (42)$$

3.2.3 Non-unimodular Lie algebras

Let us now investigate the algebraic way of looking at these Lie algebras, following MacCallum [50]. We will first investigate the ones that are non-unimodular. These are exactly those who have non-zero trace: $a_i \neq 0$. Let $\hat{a}_i$ be the unit vector parallel to a_i . For the sake of simplicity, let us assume that our frame is orientated so that $a_i = a\delta_i^4$. Then we can decompose the structure constants into

$$\begin{aligned} C_{B4}^A &= \Theta_B^A \\ n^{ab} &= \frac{1}{2} C_{ij}^a \epsilon^{ijbk} \hat{a}_k \end{aligned} \quad (43)$$

where n^{ab} is a symmetric matrix. The Jacobi identity implies $C_{ab}^4 = 0$ and thus

$$n^{a4} = n^{4a} = 0. \quad (44)$$

Note that the matrix n^{ab} only contributes to the trace-free part of the structure constants. Using a “sloppy” notation, but practical for our purposes, we can write the trace-free part as

$$\begin{aligned} D_{BC}^A &= \epsilon_{BCDk} n^{AD} \hat{a}^k \\ D_{B4}^A &= \Theta_B^A - a\delta_B^A. \end{aligned} \quad (45)$$

We have still an $SO(3)$ orientation we can use to diagonalise n^{AB} . Thus we can assume that $n^{AB} = \text{diag}(n_1, n_2, n_3)$ by choosing a suitable frame. The Jacobi identity reduces now to

$$\begin{aligned} n_1(2D_{14}^1 - a) &= n_2(2D_{24}^2 - a) = n_3(2D_{34}^3 - a) = 0 \\ n_2\Theta_2^3 + n_3\Theta_3^2 &= n_3\Theta_3^1 + n_1\Theta_1^3 = n_1\Theta_1^2 + n_2\Theta_2^1 = 0. \end{aligned} \quad (46)$$

Let us first assume that the rank of n^{ab} is 3. This leads to $D_{A4}^A = 3a/2$ which is a contradiction, since D_{bc}^a is trace-free. Hence, we can assume that $n_3 = 0$.

The classification now reduces to finding the eigenvalues n_1 and n_2 and the matrix Θ_B^A . This is done in MacCallum [50]; we will take the simplest example in this case (even though the dynamical behaviour of these types may be highly complex). Assume that the rank of n^{ab} is two. The further analysis splits in two cases, determined by the sign of $n_1 n_2$. Assume that $n_1, n_2 > 0$. This leads to $D_{14}^1 = a/2$ and $D_{24}^2 = a/2$ (and hence $D_{34}^3 = -a$). Further we get $\Theta_2^3 = \Theta_1^3 = 0$ plus the constraint $n_1\Theta_1^2 + n_2\Theta_2^1 = 0$. This algebra is type $A_{4,12}$, which MacCallum calls N22.

The other Lie algebra with $\text{rank}(n^{ab}) = 2$ is found when $n_1 < 0 < n_2$. This corresponds to the decomposable algebra $A_{2,1} \oplus A_{2,1}$. MacCallum calls this type N20.

Hence, both of these algebras, $A_{4,12}$ and $A_{2,1} \oplus A_{2,1}$, have (after choosing a orientation of frame) the following parameters:

$$n^{AB} = \text{diag}(n_1, n_2, 0), \quad D^A_{B4} = \begin{bmatrix} \frac{a}{2} & -\frac{n_1}{n_2} \Theta^2_1 & \Theta^1_3 \\ \Theta^2_1 & \frac{a}{2} & \Theta^2_3 \\ 0 & 0 & -a \end{bmatrix}. \quad (47)$$

Doing this analysis for all of the remaining cases (for rank 1 and 0) we can find the parameter space for these models. In the appendix, the general form of the matrix Θ^A_B is listed for all of the non-unimodular Lie algebras (using a particular choice of gauge). For example, in the case where $n_1 = n_2 = 0$, eq. (46) vanishes identically. Thus it remains to classify the different invariant properties of the matrix Θ^A_B . These invariant properties determine the Lie algebra type.

3.2.4 Unimodular algebras

The unimodular algebras are defined by vanishing trace: $a_i = 0$. For these algebras the following theorem holds:

Theorem (Farnsworth and Kerr) *For a four-dimensional unimodular Lie algebra there will either exist a p_a such that*

$$C^a_{bc} = \Theta^a_{[b} p_{c]}, \quad \Theta^a_b p_a = 0, \quad (48)$$

or there exists no such p_a , and there exists a non-zero ℓ^c such that

$$C^a_{bc} \ell^c = 0. \quad (49)$$

A simple and geometric proof of this is given in MacCallum [50]. MacCallum calls the algebras obeying (48) and (49), U1 and U3 respectively.

In the class U1, we can choose an orientation such that $p_a = p \delta^4_a$. Now the remaining part to be classified is the matrix Θ^A_B . Since the structure constants are trace-free, it follows that $\Theta^A_A = 0$.

The class U3 splits in two; U3I and U3S. Each of these subclasses has two members each. The members of the class U3I, $A_{4,8}$ and $A_{4,10}$, can both be seen as parametric limits of a non-unimodular model. The members of class U3S are semi-simple and are the decomposable algebras $\text{VIII} \oplus \mathbb{R}$ and $\text{IX} \oplus \mathbb{R}$.

3.3 Examples and solutions

3.3.1 The types $A_3 \oplus \mathbb{R}$

It is interesting to investigate the trivial expansion of the Bianchi types explicitly. These models correspond to the simplest Kaluza-Klein models. They have a single extra dimension, and since this dimension is homogeneous, it can be compactified into a circle.

We need to find all the structure constants possible for these models. There are two equivalent ways of doing this. Either we can do it purely algebraically, or we can do it from a geometrical point of view. We will choose the latter.

In principle, the extra dimension can be tilted. Let us choose the spatial vierbein to have three vectors spanning the vectorspace A_3 . Hence, we let three

vectors have a Bianchi-type algebra, while the fourth vector is orthogonal to A_3 . Let these three vectors which span A_3 be denoted $\mathbf{e}_B$, $B = 1, 2, 3$. We can now write

$$C_{AB}^D = \epsilon_{ABC} n^{CD} + \delta_A^D \tilde{a}_B - \delta_B^D \tilde{a}_A \quad (50)$$

with the requirement

$$\tilde{a}_D n^{CD} = 0. \quad (51)$$

We know there exists a vector $\mathbf{u}$ which commutes with $\mathbf{e}_B$. This vector is linearly independent of $\mathbf{e}_B$ so we can write $\mathbf{e}_4 = \lambda^A \mathbf{e}_A + \lambda^4 \mathbf{u}$ where $\lambda^4 \neq 0$. Hence we get

$$[\mathbf{e}_A, \mathbf{e}_4] \equiv C_{A4}^D \mathbf{e}_D = \lambda^B C_{AB}^D \mathbf{e}_D \quad (52)$$

Thus the structure constants can be written

$$C_{A4}^D = \lambda^B C_{AB}^D, \quad C_{ab}^4 = 0. \quad (53)$$

The trace of the structure constants can now be calculated

$$\begin{aligned} a_B &= \frac{1}{3} C^i_{iB} = \frac{2}{3} \tilde{a}_B \\ a_4 &= \frac{1}{3} C^i_{i4} = \frac{2}{3} \lambda^D \tilde{a}_D = \lambda^D a_D \end{aligned} \quad (54)$$

and the trace-free part is

$$\begin{aligned} D_{BC}^A &= \epsilon_{BCD} n^{DA} + \frac{1}{2} (\delta_B^A a_C - \delta_C^A a_B) \\ D_{B4}^A &= \lambda^C \epsilon_{BCD} n^{DA} - \frac{3}{2} \lambda^A a_B + \frac{1}{2} \delta_B^A a_4 \\ D_{A4}^4 &= a_A. \end{aligned} \quad (55)$$

We still have an $SO(3)$ choice of gauge to fix the orientation of the vectors $\mathbf{e}_B$, hence, we can choose a orientation where the matrix n^{CD} is diagonal.

Class A models: Let us turn our attention to the Class A models. The class A models are characterised by

$$a_B = 0.$$

The Jacobi equations and the $(0, i)$ field equations lead to a set of constraints which must be satisfied. Let us choose a frame where n^{AB} is diagonal. The constraint equations can now be satisfied with (or a frame can be chosen)

$$\begin{aligned} \sigma_{AB} &= 0 = \Omega_{AB} \quad (A \neq B) \\ \lambda_A &= 0 \\ \sigma_{4A} &= \Omega_{4A}. \end{aligned} \quad (56)$$

Note that we have not assumed $\sigma_{4A} = 0$, and in general we have not enough gauge freedom to put the shear tensor into diagonal form. The case $\mathbf{I} \oplus \mathbb{R}$ is the only case where we can diagonalise the shear vector completely. In the

other models there will still remain some off-diagonal shear components. In the absence of anisotropic stress, we will get further constraints from the off-diagonal shear equations. These can be satisfied with

$$\Omega_{4A} = 0 = \sigma_{4A}. \quad (57)$$

Thus, in this case the shear must be diagonal. Henceforth we will assume that this is the case.

The tensor b_{ab} reduces to the form

$$\begin{aligned} b_{AB} &= 2n_{AC}n^C{}_B - n^C{}_C n_{AB} + \frac{1}{2}\delta_{AB} \left[(n^C{}_C)^2 - 2n_{AC}n^C{}_B \right] \\ b_{4a} &= 0 \end{aligned} \quad (58)$$

The independent shear equations are now ($\sigma^4{}_4 = -\sigma^A{}_A$)

$$\dot{\sigma}_{AB} + \theta\sigma_{AB} + b_{AB} - \frac{1}{4}\delta_{AB}b^C{}_C = 0 \quad (59)$$

while the remaining Jacobi equations are

$$\dot{n}^{AB} + \frac{1}{4}\theta n^{AB} + \sigma^C{}_C n^{AB} - 2\sigma^{(A}{}_C n^{B)C} = 0. \quad (60)$$

Note that these are only valid in the choice of gauge mentioned above.

Example: A $\text{II} \oplus \mathbb{R}$ perfect fluid solution. Let us consider a specific case. Assume that the matrix n^{AB} has only one non-zero eigenvector. We assume that

$$n^{AB} = \text{diag}(n, 0, 0) \quad (61)$$

which corresponds to the $\text{II} \oplus \mathbb{R}$ Lie algebra. We introduce new variables $\sigma_{1,2,3}$ by

$$\begin{aligned} \sigma_{11} &= 3\sigma_1 \\ \sigma_{22} &= -\sigma_1 - \sqrt{2}\sigma_2 + \sqrt{6}\sigma_3 \\ \sigma_{33} &= -\sigma_1 - \sqrt{2}\sigma_2 - \sqrt{6}\sigma_3 \\ \sigma_{44} &= -\sigma_1 + 2\sqrt{2}\sigma_2. \end{aligned} \quad (62)$$

The equations of motion now reduce to

$$\begin{aligned} \dot{\sigma}_1 + \theta\sigma_1 + \frac{5}{24}n^2 &= 0 \\ \dot{\sigma}_2 + \theta\sigma_2 + \frac{1}{6\sqrt{6}}n^2 &= 0 \\ \dot{\sigma}_3 + \theta\sigma_3 &= 0 \\ \dot{n} + \frac{1}{4}\theta n - 5\sigma_1 n - 2\sqrt{2}\sigma_2 n &= 0. \end{aligned} \quad (63)$$

In addition to these, Raychaudhuri's equation and the energy conservation equation must be fulfilled. We will assume in this example that the perfect fluid obeys a γ -law equation of state: $p = (\gamma - 1)\rho$.

There is a specific case where we can solve the equations of motion exactly. We can search for a self-similar solution with the properties that $\sigma_{1,2,3} \propto t^{-1}$, $\rho \propto t^{-2}$ and $\theta \propto t^{-1}$. By doing this, we can find a solution which is the $\text{II} \oplus \mathbb{R}$ version of the Collins-Stewart type II perfect fluid solution. This solution is given by

$$\begin{aligned}
\sigma_1 &= -\frac{5}{66\gamma}(2\gamma-1)t^{-1} \\
\sigma_2 &= -\frac{\sqrt{2}}{33\gamma}(2\gamma-1)t^{-1} \\
\sigma_3 &= 0 \\
n^2 &= \frac{4}{11\gamma^2}(2-\gamma)(2\gamma-1)t^{-2} \\
\rho &= \frac{3}{11\gamma^2}(6-\gamma)t^{-2} \\
\theta &= \frac{2}{\gamma}t^{-1}
\end{aligned} \tag{64}$$

where $1/2 \leq \gamma \leq 2$.¹⁰ The metric can be written as

$$\begin{aligned}
ds^2 &= -dt^2 + t^{\frac{2}{11\gamma}(8-5\gamma)} \left(dx + \frac{2c}{\gamma} y dz \right)^2 \\
&\quad + t^{\frac{2}{11\gamma}(3\gamma+4)} (dy^2 + dz^2) + t^{\frac{2}{11\gamma}(6-\gamma)} dw^2
\end{aligned} \tag{65}$$

where $c^2 = (2-\gamma)(2\gamma-1)/11$. Note that if $\gamma > 8/5$ – if the matter is stiff enough – one of the dimensions will contract to an arbitrary small size.

3.3.2 Solutions to the model $A_{4,5}^{1,1}$

This model is of pure vector type and is particularly easy. We have already worked out the space of structure constants for this model. The structure constants can be characterised by a non-zero vector a_i in $\mathbb{R}^4$. We choose an orientation of the frame so that $\mathbf{e}_1$ is aligned with a_i . Hence, $a_i = \delta^1_i a$ for $a \neq 0$.

The Jacobi equations (20) and (21) are satisfied by construction, and eq. (23) is trivially satisfied. Equation (22) leads to

$$\dot{a} + \frac{1}{4}\theta a + \sigma_{11}a = 0 \tag{66}$$

and

$$\Omega_{12} = \sigma_{12}, \quad \Omega_{13} = \sigma_{13} \quad \text{and} \quad \Omega_{14} = \sigma_{14} \tag{67}$$

Eq. (29) equation tells us that

$$\sigma_{a1}a = 0 \Rightarrow \sigma_{a1} = 0. \tag{68}$$

Hence, the shear has only non-zero components in the 2, 3 and 4 directions. We still have an unused freedom of choosing the orientation of the vectors $\mathbf{e}_i$ for $i = 2, 3$ and 4. Hence, we can choose a frame where the shear σ_{ab} is diagonal.

¹⁰It should be noted that in 4+1 dimensions, radiation has $\gamma = 5/4$, and the strong energy condition requires $\gamma > 1/2$. Hence, inflation can occur for $\gamma < 1/2$.

Henceforth, we will also assume $\Lambda = 0 = \pi_{ab}$ and that the fluid obeys the barotropic equation of state $p = (\gamma - 1)\rho$.

The dynamical systems approach has proved to be a powerful tool in cosmology (see for example [53]). Let us use this method to solve the complete system of equations in the $A_{4,5}^{1,1}$ model. We parametrise the shear with

$$\sigma_{ab} = \text{diag}(0, -2\sigma_+, \sigma_+ + \sqrt{3}\sigma_-, \sigma_+ - \sqrt{3}\sigma_-) \quad (69)$$

and introduce a new time coordinate with

$$\frac{dt}{d\tau} = \frac{4}{\theta}. \quad (70)$$

Introducing the expansion normalised variables

$$\Sigma_{\pm} = \frac{2\sqrt{2}\sigma_{\pm}}{\theta}, \quad \mathcal{A} = \frac{4a}{\theta}, \quad \Omega = \frac{8\rho}{3\theta^2} \quad (71)$$

the equations of motion can be written as

$$\begin{aligned} \Sigma'_{\pm} &= (q - 3)\Sigma_{\pm} \\ \mathcal{A}' &= q\mathcal{A} \\ \Omega' &= 2[q - (2\gamma - 1)]\Omega \end{aligned} \quad (72)$$

where

$$\begin{aligned} q &= 3\Sigma + (2\gamma - 1)\Omega \\ \Sigma &= \Sigma_+^2 + \Sigma_-^2 \\ 1 &= \Sigma + \mathcal{A}^2 + \Omega. \end{aligned} \quad (73)$$

The last of these is the constraint equation, and q is the deceleration parameter defined by $\theta' = -(1 + q)\theta$.

The system of equations turns out to be quite simple, which makes it possible to solve these equations exactly. The solutions are

$$\begin{aligned} \Sigma_{\pm} &= \frac{p_{\pm}e^{-3\tau}}{(A^2e^{-6\tau} + e^{-2(2\gamma-1)\tau} + K^2)^{1/2}} \\ \Omega &= \frac{e^{-2(2\gamma-1)\tau}}{A^2e^{-6\tau} + e^{-2(2\gamma-1)\tau} + K^2} \\ \mathcal{A} &= \frac{K}{(A^2e^{-6\tau} + e^{-2(2\gamma-1)\tau} + K^2)^{1/2}} \end{aligned} \quad (74)$$

where $p_+^2 + p_-^2 = A^2$. Note that the solutions are Kasner-like near the initial singularity. At late times, the universe approaches the 5D Milne universe if $1/2 < \gamma < 2$ and the flat 5D FRW if $0 \leq \gamma < 1/2$.

We introduce the functions

$$\begin{aligned} N(t) &\equiv \left(A^2 + t^{2-\gamma} + K^2 t^{3/2}\right)^{\frac{1}{2}} \\ \beta(t) &\equiv \int \frac{Adt}{t \cdot N(t)}. \end{aligned} \quad (75)$$

The metric for the solutions can now be written

$$ds^2 = -\frac{dt^2}{N(t)^2} + t^{1/2} \left\{ dx^2 + e^{-\frac{\kappa}{2}x} \left[e^{-\sqrt{2}\cos(\phi)\beta(t)} dy^2 + e^{\sqrt{2}\cos(\phi+\pi/3)\beta(t)} dz^2 + e^{\sqrt{2}\cos(\phi-\pi/3)\beta(t)} dw^2 \right] \right\}. \quad (76)$$

3.3.3 The four-dimensional Nil^4 case

The only indecomposable Lie algebra in four dimensions that is nilpotent, is the type $A_{4,1}$. This algebra is the Lie algebra of the four-dimensional nil-geometry, denoted by Nil^4 . For the sake of illustration we will determine the possible structure constants for this model.

What we would like to have is a representation of the space $\mathcal{W}(A_{4,1})$. But finding the whole space is by no means necessary. We have a gauge freedom which corresponds to an $O(4)$ -rotation of the spatial frame. By the Gram-Schmidt process, we can show that every $A \in GL(n, \mathbb{R})$ can be written as

$$A = P \cdot R \quad (77)$$

where $R \in O(n)$ and $P \in PT(n)$ where $PT(n) \subset GL(n, \mathbb{R})$ is the group of all upper-triangular matrices with positive entries along the diagonal. Hence, by choosing a suitable frame, we can “gauge away” the rotation matrix R . It suffices therefore to look at the space

$$\mathcal{P}(\mathcal{A}) = \mathcal{W}(\mathcal{A})/O(4) = \left\{ \tilde{C}_{ij}^k | \tilde{C}_{ij}^k = (P^{-1})^k_l C^l_{nm} P^n_i P^m_j, \quad P \in PT(4) \right\}. \quad (78)$$

$PT(4)$ is a Lie group, thus if $P \in PT(4)$ then $P^{-1} \in PT(4)$. From Table 1 we have

$$C^1_{24} = 1, \quad C^2_{34} = 1. \quad (79)$$

By calculating $\mathcal{P}(\mathcal{A})$ we find that there are essentially only three non-zero commutators. These are

$$C^1_{24}, \quad C^1_{34} \quad \text{and} \quad C^2_{34}. \quad (80)$$

(see also the appendix where the $A_{4,1}$ can be seen as the $a \rightarrow 0$ limit of $A_{4,4}$.) The Lie algebra is trace-free, so $C^k_{ij} = D^k_{ij}$. Note that in the general case, the three-dimensional Ricci tensor is not diagonal. Hence, if the shear is to be diagonal, then we have to set either D^1_{24} or D^2_{34} to zero if no anisotropic stress is present. Let us assume therefore that $\pi_{ab} = 0$, $C^1_{34} = 0$, $\Omega_{ab} = 0$ and σ_{ab} diagonal.

We introduce the two curvature variables

$$N_1 = D^1_{24}, \quad N_2 = D^2_{34}. \quad (81)$$

So in this model, there will be 5 variables left to describe the geometry of the spatial hypersurfaces. These are

$$N_1, \quad N_2, \quad \sigma_{1,2,3} \quad (82)$$

where $\sigma_{1,2,3}$ are defined in eq. (62). Together with the matter equations, the evolution equation for these variables can now be written down.

In the case of a γ -law perfect fluid, one can also find a specific self-similar solution. The metric for this Nil⁴ solution is

$$ds^2 = -dt^2 + t^{\frac{2}{5}(4-3\gamma)} \left(dx + \frac{2c}{\gamma} y dw \right)^2 + t^{\frac{2}{5}(3-\gamma)} \left(dy + \frac{2c}{\gamma} z dw \right)^2 + t^{\frac{2}{5}(\gamma+2)} dz^2 + t^{\frac{2}{5}(3\gamma+1)} dw^2 \quad (83)$$

where $c^2 = (2 - \gamma)(2\gamma - 1)/10$ and $1/2 \leq \gamma \leq 2$. The matter-density of the perfect fluid is

$$\rho = \frac{3(3 - \gamma)}{5\gamma^2 t^2}. \quad (84)$$

For this solution, if $\gamma \geq 4/3$ then there will be one contracting direction. However, this solution is only a particular solution, corresponding to a fixed point in the dynamical system. The more general behaviour and the nature of this solution for the Nil⁴-world will be the subject of a future work.

3.4 Exceptional models A_4^*

For the Bianchi models in 3+1 dimensions, there is one exceptional case VI_{-1/9}^{*} for which one of the R_{0a} -equations vanish identically; it can have one extra shear degree of freedom.

In the 4+1 dimensional case, we have many such exceptional cases. By inspecting eq. (29), we see that we can have an additional shear degree of freedom in the following cases:

$$\begin{aligned} A_{4,2}^p : \quad & p = -1 \\ A_{4,5}^{pq} : \quad & q = -\frac{1+p}{2} \quad (\text{or with } q, p \text{ interchanged}) \\ A_{4,6}^{pq} : \quad & p = -q. \end{aligned}$$

We will denote these models which have this additional shear degree of freedom with A_4^* . Note that the case VI_{-1/9}^{*} $\oplus \mathbb{R}$ is obtained in the limit of $A_{4,5}^{pq*}$ as $p \rightarrow 0$:

$$\lim_{p \rightarrow 0} A_{4,5}^{pq*} = VI_{-1/9}^* \oplus \mathbb{R}. \quad (85)$$

The other exceptional models are special for the 4+1 dimensional case and have no 3+1 dimensional analogue. Note also that in the cases $A_{4,5}^{-\frac{1}{3}, -\frac{1}{3}*}$ and $A_{4,5}^{1, -1*}$ we can have *two* additional shear degrees of freedom. However, the total number of parameters remains the same because in these cases we have one less commutator function (see appendix)¹¹.

Hence, for a given Lie algebra, these exceptional models have more degrees of freedom than any of the other models.

¹¹This is because we have an additional symmetry in these cases which makes it possible to gauge away one more commutator function.

4 Compactification

As we now have presented all the homogeneous models, we note that most of the models are non-compact. The Kaluza-Klein mechanism needs a small and compact dimension to work, and hence, all of the models which do not have a compact dimension cannot be a proper model for the Kaluza-Klein mechanism. However, in many of the models we can construct compact versions of the models. In a Kaluza-Klein model one usually have a product space

$$M \times N \tag{86}$$

where M is the 3+1 dimensional base space and N is a compact space. For example, in string theory N is a Calabi-Yau manifold with 3 complex dimensions. In our model, which has in total 5 dimensions, N must be one-dimensional, and thus $N = S^1$.

However, one could instead imagine a more general model where the extra dimension is small, compact and perhaps “twisted”. Hence, we do not necessarily need to constrain ourself to a product space, but we can allow for a more general space. We can assume that our space is a *fiberbundle* P with compact fibers. For a fiberbundle there will also exist a projection map $\pi : P \mapsto M$. M is called the base space, and in our context this is our four-dimensional spacetime. We demand that for a $p \in P$ we have

$$\pi^{-1}(p) = S^1. \tag{87}$$

Locally we will not see the difference between these “twisted” spaces and the product spaces; for any $p \in P$ there will exist an open neighbourhood $U \in M$ such that

$$U \times S^1 \subset P. \tag{88}$$

The difference will be at a global scale. In a cosmological setting, the space M will be of cosmological size, and hence, we have to go to cosmological scales to see the difference between the twisted and the un-twisted versions. Notwithstanding, in the very early universe the cosmological scales may have been comparable to the small extra dimensions. Thus these twisted spaces may have had an important impact on the early evolution of the universe.

To construct a compact dimension we can proceed as follows. We find a discrete subgroup $\Gamma \subset \text{Isom}(P)$ which acts freely and properly discontinuously on P . We identify now points p and q for which there exist a $\gamma \in \Gamma$ such that $p = \gamma(q)$. There are usually many such groups for a homogeneous manifold, but not always. Also in many cases, the compactification radius can vary as we move along the base space. For the simply transitive models this compactification can always be done. These models have four linearly independent Killing vectors, ξ_i , which act freely on the spatially homogeneous hypersurfaces. Hence, for any Killing vector ξ which is a linear combination of the ξ_i ’s we can – through exponentiation $\phi = \exp \xi$ – find such a freely and properly discontinuous acting group Γ generated by the element ϕ .

In the decomposable models, the compactification depends on the single extra dimension, which is trivial. In many of these cases, we can also compactify the base space as well, and in three dimensions this has some very interesting

consequences [54, 55, 56, 57, 58, 59]. In particular, due to Mostow's rigidity theorem, all compact hyperbolic spaces of dimension three or higher, are rigid; i.e. they do not allow an anisotropic expansion. For the Bianchi types III and VIII (and hence types $\text{III} \oplus \mathbb{R}$ and $\text{VIII} \oplus \mathbb{R}$ as well) the effect of compactification is quite the opposite; they can have a unbounded number of free parameters. The compactification induces so-called moduli parameters which will increase the number of free parameters. For example, the vacuum type $\text{I} \oplus \mathbb{R}$ model has 2 free parameters, while the compactified version of type $\text{I} \oplus \mathbb{R}$ into a four-torus T^4 has 2+16 free parameters¹². This may have a significant effect on a quantum theory of gravity and on quantum cosmological models.

We would leave the compactification question open, but we will provide with an example in which we compactify both the base space and its fibers.

Compactification of the nilpotent geometries. We start by considering a three-dimensional torus T^3 . The torus can be constructed from the usual $\mathbb{E}^3$ by identification under a discrete group of translations.

Consider the product between the torus and the finite interval $[0, 1]$

$$T^3 \times [0, 1]. \quad (89)$$

The torus can be viewed upon as the unit cube in $\mathbb{E}^3$ with the usual identification of the boundary. We will now identify the two tori on boundary of (89) as follows. Take a matrix $A \in SL(3, \mathbb{Z})$ (the special linear group with integer entries). $SL(3, \mathbb{Z})$ is the mapping class group of T^3 , hence, the mapping $A(T^3)$ maps the torus isometrically onto itself. To obtain a compact space we can therefore identify the boundary $T^3 \times \{0\}$ with $T^3 \times \{1\}$ under the mapping of A . Thus if $p, q \in T^3$ then

$$\{p\} \times \{0\} \sim \{q\} \times \{1\}, \text{ iff } A(q) = p. \quad (90)$$

We have now constructed a compact manifold $M \cong T^3 \times [0, 1] / \sim$. If the matrix A is the identity matrix, then $M \cong T^4$ and hence, is of type $\text{I} \oplus \mathbb{R}$. The three other possibilities are the two nilpotent spaces¹³ $\text{Nil}^3 \times S^1$ and Nil^4 , and the solvable case $\text{Sol}^3 \times S^1$.

We define the characteristic polynomial $p_A(\lambda)$ by

$$p_A(\lambda) \equiv \det(A - \lambda I). \quad (91)$$

The roots of $p_A(\lambda)$ determine whether we have a nilpotent group or not.

We have a nilpotent geometry if all three roots of A are equal to 1. If this is not the case, then we have the solvable case $\text{Sol}^3 \times S^1$. All of the nilpotent geometries in four dimensions can be compactified completely in this way, depending on the reducibility of the matrix A .

5 Conclusion and Outlook

In this paper we have classified all spatially homogeneous cosmological models of dimension 4+1 where the spatial hypersurfaces are connected and simply

¹²For the 3+1 dimensional case this is illustrated in [60].

¹³ $\text{Nil}^3 \times S^1$ has the type $\text{II} \oplus \mathbb{R}$ while the $\text{Sol}^3 \times S^1$ has a type $VI_0 \oplus \mathbb{R}$ Lie algebra.

connected. We found five multiply transitive models which cannot be seen upon as special cases of the simply transitive models. These are in some sense the four-dimensional versions of the Kantowski-Sachs (KS) model. Three of these models were directly linked to the KS case since they consisted of products of manifolds where at least one of the components was a sphere S^2 .

For the simply transitive models, the classification of the four dimensional Lie algebras provides us with all the possible models. Among these models we should expect many interesting phenomena. For example, they may give us an understanding of how extra dimensions can affect the evolution of our four dimensional universe. The idea of extra dimensions is by no means a new idea, but higher-dimensional cosmologies seems to be little understood.

A special class of solutions to the field equations are particularly interesting. The plane-wave solutions are known to be solutions describing gravitational waves propagating through spacetime. A feature of these solutions is that they possess an extra symmetry in addition to those arising from the requirement of spatial homogeneity. These solutions have a null Killing vector. The total symmetry groups are therefore higher than the generic solution of the homogeneous field equations. These solutions also have a particular interest in string theory because they admit supersymmetry. In 4+1 dimensions there exists a large class of plane wave solutions. As a matter of fact, there is a five parameter family of vacuum plane-wave solution in 4+1 dimensions. Some of them generalises the known plane wave solutions in 3+1 dimensions, others are new and special for 4+1 dimensions. These solutions will be the subject of a future work.

A recent work which should be mentioned in relation to this, is a work by De Smet [61]. In this work, a Petrov classification of algebraically special five-dimensional spacetimes is constructed. However, what special role these spacetimes may have for multidimensional cosmology is not known.

Acknowledgments

The author is deeply grateful to J.D. Barrow for useful comments and suggestions on this work.

This work was founded by the Research Council of Norway.

Appendix: The structure constants for the indecomposable Lie algebras

We will here give a list of all the structure constants for the indecomposable ones. We have chosen a specific gauge (orientation of frame) to simplify the expressions and clarify their structure. We write them in terms of the two matrices $n^{AB} = \text{diag}(n_1, n_2, 0)$ and $C^A_{B4} = \Theta^A_B$. Note also that the obtained bound for the structure constants may in some cases not hold globally.

$$\text{rank}(n^{AB}) = 2 : n_1, n_2 \neq 0$$

$$\Theta^A_B = \begin{bmatrix} \frac{3}{2}a & -\frac{n_1}{n_2}\Theta^2_1 & \Theta^1_3 \\ \Theta^2_1 & \frac{3}{2}a & \Theta^2_3 \\ 0 & 0 & 0 \end{bmatrix}. \quad (92)$$

- $A_{4,12}$: $0 < n_1, n_2$
- $A_{2,1} \oplus A_{2,1}$: $n_1 < 0 < n_2$.

$\mathbf{rank}(n^{AB}) = 1$: $n_1 \neq 0, n_2 = 0$

$$\Theta^A_B = \begin{bmatrix} \frac{3}{2}a & \Theta^1_2 & \Theta^1_3 \\ 0 & \Theta^2_2 & \Theta^2_3 \\ 0 & \Theta^3_2 & \Theta^3_3 \end{bmatrix}. \quad (93)$$

- $A_{4,9}^q$: $\Theta^3_2 = 0, \Theta^2_2 > 0, q\Theta^2_2 = \Theta^3_3, \Theta^2_2 + \Theta^3_3 = \frac{3}{2}a$
 $q = 1 \Rightarrow \Theta^2_3 = 0$.
- $A_{4,7}$: $\Theta^3_2 = 0, \Theta^2_3 > 0, \Theta^2_2 = \Theta^3_3 > 0$
- $A_{4,11}^q$: $\text{Tr}(\mathbb{T}) = \frac{3}{2}a, 4q^2\det(\mathbb{T}) = (\text{Tr}(\mathbb{T}))^2(1 + q^2)$ where

$$\mathbb{T} = \begin{bmatrix} \Theta^2_2 & \Theta^2_3 \\ \Theta^3_2 & \Theta^3_3 \end{bmatrix} \quad (94)$$

$\mathbf{rank}(n^{AB}) = 0$: $n_1 = n_2 = 0$

$$\Theta^A_B = \begin{bmatrix} \Theta^1_1 & \Theta^1_2 & \Theta^1_3 \\ 0 & \Theta^2_2 & \Theta^2_3 \\ 0 & 0 & \Theta^3_3 \end{bmatrix}, \quad \Theta^A_A = 3a. \quad (95)$$

- $A_{4,5}^{pq}$: $\Theta^1_1 > 0, p\Theta^1_1 = \Theta^2_2, q\Theta^1_1 = \Theta^3_3$.
 $p = 1 \Rightarrow \Theta^1_2 = 0, \quad p = q \Rightarrow \Theta^2_3 = 0, \quad p = q = 1 \Rightarrow \Theta^1_3 = 0$
- $A_{4,2}^p$: $\Theta^2_2 = \Theta^3_3 \neq 0, \Theta^1_1 = p\Theta^2_2, \Theta^2_3 > 0$,
 $p = 1 \Rightarrow \Theta^1_2 = 0$.
- $A_{4,3}$: $\Theta^2_2 = \Theta^3_3 = 0, \Theta^2_3 > 0$
- $A_{4,4}$: $\Theta^1_1 = \Theta^2_2 = \Theta^3_3 = a, \Theta^2_3 > 0$
- $A_{4,6}^{pq}$: $\Theta^1_1 = 3ap/(p + 2q) \neq 0, \text{Tr}(\mathbb{T}) = 6aq/(p + 2q) > 0, 4q^2\det(\mathbb{T}) = (\text{Tr}(\mathbb{T}))^2(1 + q^2)$ where $\mathbb{T}$ is defined in eq. (94).

The rest of the non-unimodular Lie algebras can be obtained by taking various parametric limits of the above algebras.

Letting the trace go to zero, leads to

$$\begin{aligned} a &\longrightarrow 0 \\ A_{4,4} &\longrightarrow A_{4,1} \end{aligned} \quad (96)$$

and hence, all the indecomposable Lie algebras and the non-unimodular algebras can be extracted from the above. The remaining ones, the decomposable of class A, are included in the analysis in section 3.3.

References

- [1] *Modern Kaluza-Klein Theories*, eds. T. Appelquist, A. Chodos, and P.G.O. Freund, Addison-Wesley (1987)
- [2] T. Kaluza, *Sitzungsber. Preuss. Akad. Wiss. Phys. Math.* **K1** (1921) 966
- [3] O. Klein, *Z. Phys.* **37** (1926) 875
- [4] O. Klein, *Nature* **118** (1926) 516
- [5] J. Polchinski, *String Theory*, Cambridge University Press, (1998)
- [6] B. Green, *Lecture notes from TASI-96* (1997) [hep-th/9702155](#)
- [7] L. Bianchi, *Mem. Mat. Fis. Soc. It. Sc.*, Serie Terza, **XI** (1898) 267 (English translation in *Gen. Rel. Grav.* **33** (2001) 2171)
- [8] G. Fubini, *Ann. Mat. Pura Appl.* **9** (1904) 33
- [9] W.P. Thurston, *Three-Dimensional Geometry and Topology*, Princeton Uni. Press (1997)
- [10] W.P. Thurston, *Bull. Am. Math. Soc.* **6** (1982) 357
- [11] P. Forgacs and Z. Horvath, *Gen. Rel. Grav.* **10** (1979) 931
- [12] P. Forgacs and Z. Horvath, *Gen. Rel. Grav.* **11** (1979) 205
- [13] A. Chodos and S. Detweiler, *Phys. Rev.* **D21** (1980) 2167
- [14] H. Ishihara, *Prog. Theo. Phys.* **72** (1984) 376
- [15] J.D. Barrow and J. Stein-Schabes, *Phys. Rev.* **D32** (1985) 1595
- [16] J. Demaret, J.-L. Hanquin, M. Henneaux and P. Spindel, *Nucl. Phys.* **B252** (1985) 538
- [17] M. Demianski, M. Heller and M. Szydlowski, *Phys. Rev.* **D36** (1987) 2945
- [18] M. Demianski, M. Szydlowski and J. Szczesny, *Gen. Rel. Grav.* **22** (1990) 1217
- [19] A. Beloborodov, M. Demianski, P. Ivanov and A.G. Polnarev, *Phys. Rev.* **D48** (1993) 503
- [20] P. Halpern and D. Klabučar, *Gen. Rel. Grav.* **22** (1990) 1217
- [21] D.J. McManus, *Gen. Rel. Grav.* **24** (1992) 659
- [22] A.A. Coley and D.J. McManus, *J. Math. Phys.* **36** (1995) 335
- [23] G. Abolghasem, A.A. Coley and D.J. McManus, *J. Math. Phys.* **37** (1996) 361
- [24] G. Abolghasem, A.A. Coley and D.J. McManus, *Gen. Rel. Grav.* **30** (1998) 1569

- [25] S. Hervik, *Gen. Rel. Grav.* **33** (2001) 2027
- [26] P. Halpern, [gr-qc/0203055](#)
- [27] P. Halpern, [gr-qc/0207056](#)
- [28] S.M. Carroll, J. Geddes, M.B. Hoffman, and R.M. Wald, [hep-th/0110149](#)
- [29] D. Lorenz-Petzold, *Phys. Lett.* **148B** (1984) 43
- [30] D. Lorenz-Petzold, *Phys. Lett.* **149B** (1984) 79
- [31] D. Lorenz-Petzold, *Phys. Lett.* **151B** (1985) 105
- [32] D. Lorenz-Petzold, *Phys. Lett.* **158B** (1985) 110
- [33] D. Lorenz-Petzold, *Phys. Lett.* **167B** (1986) 157
- [34] D. Lorenz-Petzold, *Prog. Theo. Phys.* **73** (1985) 533
- [35] D. Lorenz-Petzold, *Class. Quant. Grav.* **2** (1985) 829
- [36] D. Lorenz-Petzold, *J. Astrophys. Astr.* **5** (1985) 137
- [37] J. Demaret and J.-L. Hanquin, *Phys. Rev.* **D31** (1985) 258
- [38] P. Halpern, *Phys. Rev.* **D33** (1986) 354
- [39] S. Ishihara, *J. Math. Soc. Japan* **7** (1955) 345
- [40] S. Kobayashi, *Transformation Groups in Differential Geometry*, Springer Verlag (1972)
- [41] W.H. Goldman, *Complex Hyperbolic Geometry*, Oxford University Press (1999)
- [42] G.F.R. Ellis and M.A.H. MacCallum, *Comm. Math. Phys.* **12** (1969) 108
- [43] G.M. Mubarakzyanov, *Izv. Vysshikh. Uchebn. Zavedenii Mat.* **1(33)** (1963) 114
- [44] G.M. Mubarakzyanov, *Izv. Vysshikh. Uchebn. Zavedenii Mat.* **3(34)** (1963) 99
- [45] G.M. Mubarakzyanov, *Izv. Vysshikh. Uchebn. Zavedenii Mat.* **4(35)** (1963) 104
- [46] S. Lie, *Theorie der Transformationsgruppen* with F. Engel, Leipzig: Teubner Verlag (1893)
- [47] J. Patera, R.T. Sharp and P. Winternitz, *J. Math. Phys.* **17** (1976) 986
- [48] J. Patera and P. Winternitz, *J. Math. Phys.* **18** (1977) 1449
- [49] G.J. Fee, *Homogeneous Spacetimes*, Master of Mathematics Thesis, Uni. of Waterloo (1979)

- [50] M.A.H. MacCallum, *On the enumeration of the real four-dimensional Lie algebras* In A.L. Harvey, editor, *On Einstein's Path: essays in honor of Engelbert Schucking*, pages 299-317. Springer Verlag, New York (1999)
- [51] C.G. Behr, *Z. Astroph.* **54** (1962) 268
- [52] A.A. Kirillov and Yu.A. Neretin, *Amer. Math. Soc. Transl.* (2) **137** (1987) 21
- [53] C.G. Hewitt and J. Wainwright in *Dynamical Systems in Cosmology*, eds: J. Wainwright and G.F.R. Ellis, Cambridge University Press (1997)
- [54] A. Ashtekar and J. Samuel, *Class. Quantum Grav.* **8** (1991) 2191
- [55] Y. Fujiwara, H. Ishihara and H. Kodama, *Class. Quantum Grav.* **10** (1993) 859
- [56] H. Kodama, *Prog. Theor. Phys.* **99** (1998) 173
- [57] J.D. Barrow and H. Kodama, *Class. Quantum Grav.* **18** (2001) 1753
- [58] J.D. Barrow and H. Kodama, *Int. J. Mod. Phys. D* **10** (2001) 785
- [59] H. Kodama, [gr-qc/0109064](#)
- [60] S. Hervik, *Class. Quantum Grav.* **17** (2000) 2765
- [61] P.-J. De Smet, [hep-th/0206106](#)